



**Letter of Intent:**  
**A New QCD facility at the M2 beam line of the CERN SPS\***  
**COMPASS++<sup>†</sup>/AMBER<sup>‡</sup>**

B. Adams<sup>13,12</sup>, C.A. Aidala<sup>1</sup>, R. Akhunzyanov<sup>14</sup>, G.D. Alexeev<sup>14</sup>, M.G. Alexeev<sup>41</sup>, A. Amoroso<sup>41,42</sup>,  
 V. Andrieux<sup>44</sup>, N.V. Anfimov<sup>14</sup>, V. Anosov<sup>14</sup>, A. Antoshkin<sup>14</sup>, K. Augsten<sup>14,32</sup>, W. Augustyniak<sup>46</sup>,  
 C.D.R. Azevedo<sup>4</sup>, A. Azhibekov<sup>2</sup>, B. Badelek<sup>47</sup>, F. Balestra<sup>41,42</sup>, M. Ball<sup>8</sup>, J. Barth<sup>9</sup>, R. Beck<sup>8</sup>,  
 Y. Bedfer<sup>20</sup>, J. Berenguer Antequera<sup>41,42</sup>, J.C. Bernauer<sup>34,45</sup>, J. Bernhard<sup>19</sup>, M. Bodlak<sup>31</sup>, P. Bordalo<sup>23,a</sup>,  
 F. Bradamante<sup>39</sup>, A. Bressan<sup>38,39</sup>, M. Büchele<sup>16</sup>, V.E. Burtsev<sup>40</sup>, W.-C. Chang<sup>35</sup>, C. Chatterjee<sup>11</sup>,  
 X. Chen<sup>21</sup>, M. Chiosso<sup>41,42</sup>, A.G. Chumakov<sup>40</sup>, S.-U. Chung<sup>17,b</sup>, A. Cicuttin<sup>39,c</sup>, P. Correia<sup>4</sup>,  
 M.L. Crespo<sup>39,c</sup>, S. Dalla Torre<sup>39</sup>, S.S. Dasgupta<sup>11</sup>, S. Dasgupta<sup>38,39</sup>, N. Dashyan<sup>50</sup>, I. Denisenko<sup>14</sup>,  
 O.Yu. Denisov<sup>42</sup>, L. Dhara<sup>11</sup>, N. d'Hose<sup>20</sup>, F. Donato<sup>43</sup>, S.V. Donskov<sup>33</sup>, N. Doshita<sup>49</sup>,  
 Ch. Dreisbach<sup>17</sup>, W. Dünnweber<sup>d</sup>, R.R. Dusaev<sup>40</sup>, M. Dziewiecki<sup>48</sup>, A. Dzyuba<sup>18</sup>, A. Efremov<sup>14</sup>,  
 P. Egelhof<sup>15</sup>, A. Elagin<sup>13</sup>, P.D. Eversheim<sup>8</sup>, M. Faessler<sup>d</sup>, J. Fedotova<sup>26</sup>, A. Ferrero<sup>20</sup>, M. Finger<sup>31</sup>,  
 M. Finger jr.<sup>31</sup>, H. Fischer<sup>16</sup>, C. Franco<sup>23</sup>, J.M. Friedrich<sup>17</sup>, V. Frolov<sup>14</sup>, A. Futch<sup>44</sup>, F. Gautheron<sup>44</sup>,  
 O.P. Gavrichtchouk<sup>14</sup>, S. Gerassimov<sup>28,17</sup>, S. Gevorkyan<sup>14</sup>, Y. Ghandilyan<sup>50</sup>, J. Giarra<sup>25</sup>, I. Gnesi<sup>41,42</sup>,  
 M. Gorzellik<sup>16</sup>, A. Grasso<sup>41,42</sup>, A. Gridin<sup>14</sup>, M. Grosse Perdekamp<sup>44</sup>, B. Grube<sup>17</sup>, R.I. Gushterski<sup>14</sup>,  
 A. Guskov<sup>14</sup>, D. Hahne<sup>9</sup>, G. Hamar<sup>39</sup>, D. von Harrach<sup>25</sup>, X. He<sup>3</sup>, R. Heitz<sup>44</sup>, F. Herrmann<sup>16</sup>,  
 N. Horikawa<sup>30,e</sup>, C.-Y. Hsieh<sup>35,f</sup>, S. Huber<sup>17</sup>, A. Inglessi<sup>18</sup>, A. Ilyichev<sup>26</sup>, T. Isataev<sup>2</sup>, S. Ishimoto<sup>49,g</sup>,  
 A. Ivanov<sup>14</sup>, N. Ivanov<sup>50</sup>, Yu. Ivanshin<sup>14</sup>, T. Iwata<sup>49</sup>, M. Jandek<sup>31</sup>, V. Jary<sup>32</sup>, C.-M. Jen<sup>24</sup>, R. Joosten<sup>8</sup>,  
 P. Jörg<sup>16</sup>, K. Juraskova<sup>32</sup>, E. Kabuß<sup>25</sup>, A. Kabyshev<sup>2</sup>, F. Kaspar<sup>17</sup>, A. Kerbizi<sup>38,39</sup>, B. Ketzer<sup>8</sup>,  
 G.V. Khaustov<sup>33</sup>, Yu.A. Khokhlov<sup>33,h</sup>, M. Kim<sup>1</sup>, O. Kiselev<sup>15</sup>, Yu. Kisselev<sup>14</sup>, F. Klein<sup>9</sup>,  
 J.H. Koivuniemi<sup>44</sup>, V.N. Kolosov<sup>33</sup>, K. Kondo<sup>49</sup>, I. Konorov<sup>28,17</sup>, V.F. Konstantinov<sup>33</sup>,  
 A.M. Kotzinian<sup>42,j</sup>, O.M. Kouznetsov<sup>14</sup>, A. Koval<sup>46</sup>, Z. Kral<sup>32</sup>, F. Krinner<sup>17</sup>, Y. Kulinich<sup>44</sup>, F. Kunne<sup>20</sup>,  
 K. Kurek<sup>46</sup>, R.P. Kurjata<sup>48</sup>, K. Kuterbekov<sup>2</sup>, I.I. Kuznetsov<sup>40</sup>, A. Kveton<sup>32</sup>, E.A. Levchenko<sup>40</sup>,  
 S. Levorato<sup>39</sup>, Y.-S. Lian<sup>35,k</sup>, Y. Liang<sup>21</sup>, J. Lichtenstadt<sup>36</sup>, P.-J. Lin<sup>20</sup>, K. Liu<sup>24</sup>, M.X. Liu<sup>24</sup>,

\* email: NQF-M2@cern.ch

<sup>†</sup> Common Muon Proton Apparatus for Structure and Spectroscopy

<sup>‡</sup> Apparatus for Meson and Baryon Experimental Research

R. Longo<sup>44</sup>, W. Lorenzon<sup>1</sup>, M. Losekamm<sup>17</sup>, V.E. Lyubovitskij<sup>40,1</sup>, E. Maev<sup>18</sup>, A. Maggiora<sup>42</sup>,  
A. Magnon, V. Makarenko<sup>26</sup>, N. Makins<sup>44</sup>, N. Makke<sup>39,c</sup>, G.K. Mallot<sup>19</sup>, A. Maltsev<sup>14</sup>, S.A. Mamon<sup>40</sup>,  
B. Marianski<sup>46</sup>, A. Martin<sup>38,39</sup>, H. Marukyan<sup>50</sup>, J. Marzec<sup>48</sup>, N. Masi<sup>7</sup>, J. Matoušek<sup>38,39</sup>, H. Matsuda<sup>49</sup>,  
G. Mattson<sup>44</sup>, G.V. Meshcheryakov<sup>14</sup>, W. Meyer<sup>6</sup>, M. Meyer<sup>44,20</sup>, Yu.V. Mikhailov<sup>33</sup>, M. Mikhasenko<sup>8</sup>,  
M. Minot<sup>13,12</sup>, E. Mitrofanov<sup>14</sup>, Y. Miyachi<sup>49</sup>, A. Moretti<sup>38</sup>, A. Nagaytsev<sup>14</sup>, C. Naïm<sup>20</sup>, D. Neyret<sup>20</sup>,  
G. Nigmatkulov<sup>29</sup>, A. Mkrtchyan<sup>50</sup>, H. Mkrtchyan<sup>50</sup>, J. Nový<sup>32</sup>, W.-D. Nowak<sup>25</sup>, F. Nozzoli<sup>37</sup>,  
G. Nukazuka<sup>49</sup>, A.S. Nunes<sup>23</sup>, A.G. Olshevsky<sup>14</sup>, I. Orlov<sup>14</sup>, J.D. Osborn<sup>1</sup>, M. Ostrick<sup>25</sup>,  
D. Panzieri<sup>42,m</sup>, B. Parsamyan<sup>41,42</sup>, S. Paul<sup>17</sup>, J.-C. Peng<sup>44</sup>, F. Pereira<sup>4</sup>, D.V. Peshekhonov<sup>14</sup>,  
M. Pešek<sup>31</sup>, M. Pešková<sup>31</sup>, N. Pierre<sup>25,20</sup>, S. Platchkov<sup>20</sup>, J. Pochodzalla<sup>25</sup>, V.A. Polyakov<sup>33</sup>,  
T. Pöschl<sup>17</sup>, J. Pretz<sup>9,i</sup>, M. Quaresma<sup>35</sup>, C. Quintans<sup>23</sup>, S. Ramos<sup>23,a</sup>, R. Raymond<sup>1</sup>, G. Reicherz<sup>6</sup>,  
C. Riedl<sup>44</sup>, D.I. Ryabchikov<sup>33,17</sup>, A. Rybnikov<sup>14</sup>, A. Rychter<sup>48</sup>, R. Salac<sup>32</sup>, V.D. Samoylenko<sup>33</sup>,  
A. Sandacz<sup>46</sup>, S. Sarkar<sup>11</sup>, M. Sarsour<sup>3</sup>, I.A. Savin<sup>14</sup>, G. Sbrizzai<sup>38,39</sup>, H. Schmieden<sup>9</sup>, E. Seder<sup>20</sup>,  
A. Selyunin<sup>14</sup>, H. Simon<sup>15</sup>, L. Sinha<sup>11</sup>, S. Sirtl<sup>16</sup>, M. Slunicka<sup>14</sup>, J. Smolik<sup>14</sup>, A. Srnka<sup>10</sup>, D. Steffen<sup>17</sup>,  
M. Stolarski<sup>23</sup>, O. Subrt<sup>32</sup>, M. Sulc<sup>22</sup>, H. Suzuki<sup>49,e</sup>, A. Szabelski<sup>38,39</sup>, P. Sznajder<sup>46</sup>, M. Tasevsky<sup>14</sup>,  
V. Tchekhovskii<sup>26</sup>, S. Tessaro<sup>39</sup>, F. Tessarotto<sup>39</sup>, A. Thiel<sup>8</sup>, J. Tomsa<sup>31</sup>, F. Tosello<sup>42</sup>, V. Tskhay<sup>28</sup>,  
S. Uhl<sup>17</sup>, B.I. Vasilishin<sup>40</sup>, A. Vassiliev<sup>18</sup>, A. Vauth<sup>9</sup>, B.M. Veit<sup>19,25</sup>, B. Ventura<sup>20</sup>, J. Veloso<sup>4</sup>,  
A. Vidon<sup>20</sup>, M. Virius<sup>32</sup>, A. Vorobyev<sup>18</sup>, M. Wagner<sup>8</sup>, S. Wallner<sup>17</sup>, M. Wilfert<sup>25</sup>, Z. Xiao<sup>5</sup>, N. Xu<sup>21</sup>,  
J. Yang<sup>21</sup>, K. Zaremba<sup>48</sup>, P. Zavada<sup>14</sup>, M. Zavertyaev<sup>28</sup>, E. Zemlyanichkina<sup>14</sup>, H. Zhao<sup>21</sup>, Y. Zhao<sup>39</sup>,  
N. Zhuravlev<sup>14</sup>, M. Ziembicki<sup>48</sup>, P. Zuccon<sup>37</sup>

<sup>1</sup> Randall Laboratory of Physics, University of Michigan, Ann Arbor, MI 48109-1040, USA

<sup>2</sup> L.N.Gumilyov Eurasian National University, Astana, Kazakhstan

<sup>3</sup> Georgia State University, Atlanta, GA 30302, USA

<sup>4</sup> University of Aveiro, i3N, Dept. of Physics, 3810-193 Aveiro, Portugal

<sup>5</sup> Tsinghua University, Beijing, 100084, P. R. China

<sup>6</sup> Universität Bochum, Institut für Experimentalphysik, 44780 Bochum, Germany

<sup>7</sup> INFN Sezione di Bologna, Bologna, Italy

<sup>8</sup> Universität Bonn, Helmholtz-Institut für Strahlen- und Kernphysik, 53115 Bonn, Germany

<sup>9</sup> Universität Bonn, Physikalisches Institut, 53115 Bonn, Germany

<sup>10</sup> Institute of Scientific Instruments, AS CR, 61264 Brno, Czech Republic

<sup>11</sup> Matrivani Institute of Experimental Research & Education, Calcutta-700 030, India

<sup>12</sup> Incom Inc., Charlton, MA 01507, USA

<sup>13</sup> Enrico Fermi Institute, University of Chicago, Chicago, IL 60637, USA

<sup>14</sup> Joint Institute for Nuclear Research, 141980 Dubna, Moscow region, Russia

<sup>15</sup> GSI Helmholtzzentrum für Schwerionenforschung, 64291 Darmstadt, Germany

<sup>16</sup> Universität Freiburg, Physikalisches Institut, 79104 Freiburg, Germany

<sup>17</sup> Technische Universität München, Physik Dept., 85748 Garching, Germany

<sup>18</sup> Petersburg Nuclear Physics Institute, 188300 Gatchina, Russia

<sup>19</sup> CERN, 1211 Geneva 23, Switzerland

<sup>20</sup> IRFU, CEA, Université Paris-Saclay, 91191 Gif-sur-Yvette, France

<sup>21</sup> Institute of Modern Physics, Chinese Academy of Science, Lanzhou 730000, China

<sup>22</sup> Technical University in Liberec, 46117 Liberec, Czech Republic

<sup>23</sup> LIP, 1649-003 Lisbon, Portugal

<sup>24</sup> Physics Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA

<sup>25</sup> Universität Mainz, Institut für Kernphysik, 55099 Mainz, Germany

<sup>26</sup> Institute for Nuclear Problems of Belarusian State University, Minsk, Belarus

<sup>27</sup> University of Miyazaki, Miyazaki 889-2192, Japan

<sup>28</sup> Lebedev Physical Institute, 119991 Moscow, Russia

<sup>29</sup> National Research Nuclear University MEPhI, Moscow, Russia

<sup>30</sup> Nagoya University, 464 Nagoya, Japan

- <sup>31</sup> Charles University in Prague, Faculty of Mathematics and Physics, 18000 Prague, Czech Republic  
<sup>32</sup> Czech Technical University in Prague, 16636 Prague, Czech Republic  
<sup>33</sup> State Scientific Center Institute for High Energy Physics of National Research Center ‘Kurchatov Institute’, 142281 Protvino, Russia  
<sup>34</sup> Stony Brook University, SUNY, Stony Brook, NY 11794-3800, USA  
<sup>35</sup> Academia Sinica, Institute of Physics, Taipei 11529, Taiwan  
<sup>36</sup> Tel Aviv University, School of Physics and Astronomy, 69978 Tel Aviv, Israel  
<sup>37</sup> Università degli Studi di Trento and INFN-TIFPA, Trento, Italy  
<sup>38</sup> University of Trieste, Dept. of Physics, 34127 Trieste, Italy  
<sup>39</sup> Trieste Section of INFN, 34127 Trieste, Italy  
<sup>40</sup> Tomsk Polytechnic University, 634050 Tomsk, Russia  
<sup>41</sup> University of Turin, Dept. of Physics, 10125 Turin, Italy  
<sup>42</sup> Torino Section of INFN, 10125 Turin, Italy  
<sup>43</sup> Università di Torino and INFN-Torino, Turin, Italy  
<sup>44</sup> University of Illinois at Urbana-Champaign, Dept. of Physics, Urbana, IL 61801-3080, USA  
<sup>45</sup> RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, NY 11973-5000, USA  
<sup>46</sup> National Centre for Nuclear Research, 00-681 Warsaw, Poland  
<sup>47</sup> University of Warsaw, Faculty of Physics, 02-093 Warsaw, Poland  
<sup>48</sup> Warsaw University of Technology, Institute of Radioelectronics, 00-665 Warsaw, Poland  
<sup>49</sup> Yamagata University, Yamagata 992-8510, Japan  
<sup>50</sup> A.Alikhanyan National Science Laboratory, Yerevan Physics Institute Foundation, Alikhanian Br. Street, 0036, Yerevan, Armenia

<sup>a</sup> Also at Instituto Superior Técnico, Universidade de Lisboa, Lisbon, Portugal

<sup>b</sup> Also at Physics Dept., Brookhaven National Laboratory, Upton, NY 11973, USA

<sup>c</sup> Also at Abdus Salam ICTP, 34151 Trieste, Italy

<sup>d</sup> Supported by the DFG cluster of excellence ‘Origin and Structure of the Universe’ ([www.universe-cluster.de](http://www.universe-cluster.de)) (Germany)

<sup>e</sup> Also at Chubu University, Kasugai, Aichi 487-8501, Japan

<sup>f</sup> Also at Dept. of Physics, National Central University, 300 Jhongda Road, Jhongli 32001, Taiwan

<sup>g</sup> Also at KEK, 1-1 Oho, Tsukuba, Ibaraki 305-0801, Japan

<sup>h</sup> Also at Moscow Institute of Physics and Technology, Moscow Region, 141700, Russia

<sup>i</sup> Present address: RWTH Aachen University, III. Physikalisches Institut, 52056 Aachen, Germany

<sup>j</sup> Also at Yerevan Physics Institute, Alikhanian Br. Street, Yerevan, Armenia, 0036

<sup>k</sup> Also at Dept. of Physics, National Kaohsiung Normal University, Kaohsiung County 824, Taiwan

<sup>l</sup> Also at Institut für Theoretische Physik, Universität Tübingen, 72076 Tübingen, Germany

<sup>m</sup> Also at University of Eastern Piedmont, 15100 Alessandria, Italy

## Executive Summary

In this Letter of Intent, we propose a broad experimental programme for the “New QCD facility at the M2 beam line of the CERN SPS”. This unrivalled installation will provide the site for a great variety of measurements to address fundamental issues of Quantum Chromodynamics, which are expected to lead to significant improvements in the understanding of QCD as our present theory of strong interactions. The proposed measurements cover the range from lowest- $Q^2$  physics as the determination of the proton radius by elastic muon-proton scattering, over average- $Q^2$ -reactions to study hadron spectroscopy, to high- $Q^2$  hadron-structure investigations using the Drell-Yan process and Deeply Virtual Compton Scattering.

In this Executive Summary, we briefly describe a number of unique first-generation experiments that are already possible using the existing M2 beam line with muons or unseparated hadrons. We also describe a number of unique second-generation experiments that rely on high-energy high-intensity radio-frequency (RF) separated hadron beams, which could be made possible by a major upgrade of the M2 beam line. Such an upgrade, which is presently under study by CERN EN-EA in the context of the Physics beyond Colliders Initiative, would make the CERN SPS M2 beam line absolutely unique in the world for many years to come. For every proposed experiment, individual instrumentation using modern detector architecture will be constructed and installed in the experimental hall EHN2, where the upgraded multi-purpose two-stage magnetic spectrometer will serve as experimental backbone of the new facility.

### Proton radius measurement using muon-proton elastic scattering

The aim of this experiment is an independent precision determination of the electric mean-square charge radius of the proton using elastic muon-proton scattering. Such a measurement appears timely, since in spite of many years of intense activity the proton-radius puzzle remains unsolved up to now. Presently, a discrepancy as large as 5 standard deviations exists between the two most recent precision measurements:  $r_{CREMA}^{rms} = 0.841 \pm 0.001$  from line-splitting measurements in laser spectroscopy of muonic hydrogen and  $r_{MAMI}^{rms} = 0.879 \pm 0.008$  from elastic electron-proton scattering.

We propose to perform a one-year measurement using high-energy muons of the CERN M2 beam line, which will provide a new and completely independent result on the proton radius with a statistical accuracy of 0.01 and considerably smaller systematic uncertainty. Using muons instead of electrons is highly advantageous, as several experimental systematic effects and also theoretical (radiative) corrections are considerably smaller. The measurement will employ a time-projection chamber filled with pure hydrogen up to pressures of 20 bar, which serves at the same time as a target and as detector gas. It has been developed by the PNPI group for several other experiments and will be adapted in a common effort to the proton-radius measurement.

### Hard exclusive reactions using a muon beam and a transversely polarised target

The main goal of this experiment is to extend our fundamental knowledge on the angular-momentum structure of the nucleon, which can be accomplished by measurements of Generalized Parton Distributions (GPDs). These functions describe the correlations between longitudinal momentum fractions and transverse spatial positions of partons in the nucleon, commonly known as “3-dimensional” picture of the nucleon. Experimental data on these GPDs are required over the largest possible domain of the Bjorken- $x$  variable, as the evaluation of the average total angular momentum  $J^f$  carried by quarks of flavour  $f$  through the Ji sum rule requires an integration over the sum of the GPDs  $H^f$  and  $E^f$ . While some experimental knowledge on the GPDs  $H^f$  exists already for  $u$  and  $d$  quarks, no experimental data exists on the “elusive” GPDs  $E^f$ .

Access to the GPDs  $E^f$  is possible by measuring certain cross section asymmetries in Deeply Virtual Compton Scattering, using a polarised charged-lepton beam and a transversely polarised target. A measurement in the so-far uncharted Bjorken- $x$  domain between 0.005 and 0.05, where phenomenological models predict a particularly large sensitivity to  $E^f$ , can only be performed using the high-energy self-polarised muon beam of the CERN SPS M2 beam line. A 2-year measurement will yield an unprece-

dedented statistical accuracy of 0.03 for the cross section asymmetry when using four bins per variable, with the systematic uncertainty expected to be smaller. A new 3-layer Silicon detector has to be constructed and installed inside of the existing polarised target, in order to accomplish identification of the target-recoil proton by the  $dE/dx$  method and to measure its momentum and trajectory. The proposed measurement is complementary in kinematics to a JLab experiment planned for 2022, which will use a transversely-polarised HD ice target.

### **Drell-Yan and charmonium production**

The main objective of the proposed Drell-Yan and  $J/\psi$  production experiments is to make a major step forward in the determination of the nearly unknown pion and kaon parton distribution functions (PDFs). The planned measurements will provide key benchmarks for testing the most recent predictions of fundamental, non-perturbative QCD calculations, such as lattice QCD and Dyson-Schwinger Equations formalism. At medium and large values of Bjorken- $x$ , a quantitative comparison between the pion and the kaon valence distributions is of utmost importance. At smaller values of Bjorken- $x$ , improved knowledge of the onset of the sea and gluon distributions in the meson will help in explaining the differences between the gluon contents of pions, kaons and nucleons, and hopefully provide clues to understand the mechanism that generates the hadron masses. Furthermore, a comparison between cross sections for positive and negative meson beams is expected to provide stringent experimental constraints on the  $J/\psi$  production mechanism, as well as an alternative way of accessing both quark and gluon distributions in the incoming meson. In parallel to meson structure measurements, the availability of heavier nuclear targets in the setup will allow the study of cold nuclear effects such as nuclear PDFs and partonic energy loss.

The M2 secondary hadron beam line at the CERN SPS provides an exclusive opportunity for such measurements. For both Drell-Yan and  $J/\psi$  production processes, the expected statistical accuracies correspond to a few  $10^4$  and nearly  $10^6$  events for the Drell-Yan and  $J/\psi$  production processes, respectively. The first part of this programme can be completed using the presently available positive ( $\pi^+$ ,  $K^+$ ,  $p$ ) and negative ( $\pi^-$ ,  $K^-$ ,  $\bar{p}$ ) hadron beams, in combination with an optimized experimental setup. A detailed study of the presently uncharted kaon structure, which is envisaged in a later stage of the programme, requires the existence of a high-intensity kaon beam that can only be produced using radio-frequency beam-separation cavities. The proposed studies need a high-performance beam particle identification. They would also greatly benefit from a novel and compact lepton absorber/detector immersed in a magnetic field. Such detector could largely improve the performances of the setup, while ensuring a large geometric acceptance and allowing for  $e^+/e^-$  detection in parallel with the muon pair.

### **Measurement of antiproton production cross sections for Dark Matter Search**

The purpose of this experiment is the measurement of the antiproton production cross section on proton and  ${}^4\text{He}$  targets for projectile energies from several ten to a few hundred GeV. In combination with similar measurements by LHCb in the TeV range, our data will provide a fundamental data set that will greatly improve the accuracy of the predicted natural flux of antiprotons in the galactic cosmic rays. This is of great importance as the indirect detection of Dark Matter (DM) is based on the search for products of DM annihilation or decay, which are expected to appear as distortions in the spectra of rare Cosmic Ray components like positrons or antiprotons. The new data set is expected to vastly improve the sensitivity of the very accurate AMS antiproton flux measurements to DM signals, which is presently spoiled by the poor knowledge of the antiproton production cross section.

The existing M2 beam line with its momentum range between 20 – 30 and 280 GeV/c is an ideal tool to perform this measurement. The double-differential antiproton production cross section will be measured in  $p + p$  and  $p + {}^4\text{He}$  collisions using the existing spectrometer in EHN2 equipped with liquid hydrogen and liquid helium targets. Measuring for several beam momenta the cross section in 20 bins each for antiproton momentum and pseudorapidity, a statistical uncertainty of 1% will be reached, with an anticipated point-to-point systematic uncertainty of less than 5%. The main challenges for these measurements

are the precise determination of the efficiencies of the trigger systems and of the particle identification efficiency and purity.

### **Spectroscopy with low-energy antiprotons**

The main objective of this experiment is a significant improvement in the understanding of the nature of the so-called  $X$ ,  $Y$ ,  $Z$  states, which remains so far unclear in spite of worldwide efforts in the last 15 years. The discovery of narrow resonance-like signals at  $e^+e^-$  colliders, which have masses above the open-charm threshold and decay to charmonium, had triggered a wealth of theoretical interpretations ranging from conventional excited charmonium to multi-quark states and hybrid mesons. Of particular interest is the production of states in association with a recoil particle, which opens the possibility of observing states with spin-exotic quantum numbers such as hybrids or glueballs. The annihilation of low-energy antiprotons with momenta below 20 GeV provides a unique tool to study these states.

An early experiment at the M2 beamline of the SPS using low-energy antiprotons and a liquid hydrogen target can reach luminosities of the order of  $10^{30} \text{ cm}^{-2} \text{ s}^{-1}$ . It is expected to make important contributions to the production and spectroscopy of these long sought-after states even before the start of PANDA at FAIR, where such a measurement is in the planning state. Our measurements will at the same time provide vital input for PANDA, e.g. by measuring production cross sections for  $X$ ,  $Y$ ,  $Z$  states in  $\bar{p}p$  reactions, for which theoretical estimates range from 0.1 nb to 10 nb. Owing to the low beam momentum, the experiment requires a dedicated target spectrometer including charged-particle tracking and electromagnetic calorimetry, in addition to the forward spectrometer in EHN2.

### **Spectroscopy of kaons**

The goal of the kaon spectroscopy programme is to map out the complete spectrum of excited kaons with unprecedented precision using novel analysis methods. A more precise knowledge of the kaon spectrum will have a broad impact not only on low-energy QCD, but also on many processes in modern hadron and particle physics where excited kaons appear, e.g. in the study of CP violation in heavy-meson decays, which are studied at LHCb and Belle II.

The high-intensity RF-separated kaon beam of the upgraded M2 beam line is an unrivalled place to accumulate a large data set for kaon spectroscopy as most of the planned or proposed measurements in the strange-meson sector elsewhere can either not compete with the measurement proposed here or are complementary to it. Taking data over one year with a beam energy of at least 50 GeV, about  $20 \times 10^6$  events can be collected, which allows us to keep systematic effects under control. The main experimental challenges are high-precision vertex reconstruction, photon detection with electromagnetic calorimeters for reconstructing neutral hadrons, and final-state particle identification, i.e. kaons have to be distinguished from pions with high efficiency over a broad kinematic range.

### **Study of the gluon distribution in the kaon via prompt-photon production**

The purpose of this experiment is the study of the gluon content of charged kaons. This is of fundamental importance for understanding the internal structure of light mesons and may also shed light onto the mechanism that generates their mass. In contrast to the well-known gluon distribution of the nucleon, our knowledge on the gluon distribution in light mesons is rather limited. It can be considerably improved by studying prompt-photon production in hadronic collisions using a high-energy meson beam.

The high-intensity RF-separated kaon beam of the upgraded M2 beam line is an unrivalled site to investigate the gluon content of the kaon. Using a beam of positive kaons with an energy of at least 100 GeV over 1-2 years, the gluon PDF in the charged kaon will be measured using the dominant hard gluon Compton scattering process in the range of  $x_g$  above 0.05 and for  $Q^2 \sim p_T^2 > 7 (\text{GeV}/c)^2$ . A separate data set will have to be taken with a beam of negative kaons in order to allow for a separation of the subdominant quark-antiquark annihilation process. For systematic studies and also to improve the knowledge on the gluon structure of the pion, it is foreseen to also collect data with incoming pions. This can be done either in parallel or separately in a preceding running period. The basic requirements for the proposed



measurement are a sufficiently high transparency of the experimental setup for the produced photons, a wide kinematic range of photon detection by the system of electromagnetic calorimeters, efficient beam hadron identification, and a dedicated calorimeter-based trigger on high- $p_T$  photons. We are not aware of plans to study the gluon structure of charged kaons at other laboratories.

### Low-energy tests of QCD using Primakoff reactions

The main objective of this experiment is the first measurement of the kaon polarisability. Electric and magnetic polarisabilities, which describe the rigidity against deformation by an external electromagnetic field, constitute fundamental quantities of low-energy QCD. Apart from a quite vague experimental upper limit, presently only model predictions exist for the electric polarisability of the kaon, e.g. by chiral perturbation theory ( $\chi PT$ ) or the quark-confinement model. It can be determined using the so-called Primakoff reaction,  $K^- Z \rightarrow K^- \gamma Z$ , using a high-energy high-intensity kaon beam, on a nuclear target  $Z$  with  $Z$  protons. Such an experimental approach allows the parallel determination of another fundamental quantity of low-energy QCD, i.e. the lifetime of the neutral pion, which is related to the chiral-anomaly hypothesis.

The RF-separated beam of the upgraded M2 beam line is a unique place for these measurements. One year of data taking with a 100 GeV beam delivering  $5 \times 10^6$  kaons per second will allow to determine the electric kaon polarisability with a statistical accuracy of  $0.03 \times 10^{-4} \text{fm}^3$ , with the total systematic uncertainty estimated to be smaller. This accuracy is even smaller than that of the prediction of  $\chi PT$  in the one-loop approximation and will clearly allow us to distinguish between models for the kaon polarisability. The neutral pion lifetime will be measured in parallel using those beam pions that remain in the RF-separated kaon beam. Here, the resulting statistical accuracy of about 1% will be smaller by more than a factor of two compared to the accuracy of the theoretical prediction and that of existing measurements. The proposed measurement requires advanced beam particle identification, excellent tracking capabilities at small angles and precise electromagnetic calorimeters. All these expectations are estimated to be met by the upgraded spectrometer that is planned to be available in EHN2 at the time of running. The proposed measurements have presently no competitors in the world.

### Production of vector mesons and excited kaons off nuclei

The main goal of this measurement is to study pion-induced exclusive vector-meson production off various nuclei in order to determine the nuclear dependence of the cross section for longitudinally polarised vector mesons ( $\rho^0(770)$ ,  $K^{0*}(892)$ ). The absorption strength for a vector meson produced in a nuclear environment is predicted to strongly depend on its polarisation and also on the beam energy. However, neither existing data nor phenomenological models allow presently for a satisfactory description of nuclear transparency. Of particular interest is a clarification of a possible colour screening effect, as predicted by the colour transparency model. When using incoming kaons, the absorption of polarised  $K^*$  mesons can be studied, which is a topic of increasing interest.

About one year of running using pions and kaons produced by the M2 beam line is required to reach a sufficient accuracy of the cross section measurement. The beam energy should be higher than about 50 GeV to avoid effects related to meson decays in the nuclear medium and lower than about 100 GeV as the cross section for the charge-exchange reaction decreases with energy. In this energy range, the cross section is sufficiently large, so that parallel running with other physics programmes can be considered. The proposed measurements are kinematically fully complementary to the studies that are planned to be performed at JLab.

### Tentative schedule of the CERN M2 beam line

A tentative schedule of the CERN M2 beam line is outlined in table 1 according to the information given on <https://lhc-commissioning.web.cern.ch/lhc-commissioning/schedule/LHC-long-term.htm>.

<b>Year</b>	<b>Activity</b>	<b>Duration</b>	<b>Beam</b>
2019 2020	Long Shutdown 2	2 years	-
2021	COMPASS-II transversity with polarised deuteron target	1 year	muon
2022	proton radius	1 year	muon
2023 2024	Drell-Yan for $\pi$ and $K$ PDFs and charmonium production mechanism	$\lesssim 2$ years	$p, K^+, \pi^+$ $\bar{p}, K^-, \pi^-$
	Antiproton cross section for Dark Matter Search	2 month	$p$
2025	Long Shutdown 3 (for SPS)		

Table 1: Tentative schedule of the CERN M2 beam line for the time between LS2 and LS3.